



At last, a scabies therapy you can trust with your smallest patients.³

Learn more about FDA-approved Natroba™ for your practice.


(spinosad) Topical Suspension, 0.9%

Children deserve better.

Twenty-nine percent of all scabies prescriptions in the US are for children under the age of 18 years.¹ The need for an agent with a better pediatric safety profile is long overdue.

29% of all scabies prescriptions in the US are for children under the age of 18.

Natroba™ is now available.²⁻⁵

- No evidence of systemic absorption of the active compound, spinosad
- No neurotoxicity of the active compound, spinosad
- No known resistance challenges associated with current agents used in today's treatment of scabies



Ensure your patients have the opportunity to achieve a “complete cure.” Initiate treatment with Natroba™.³

Learn more about the safe, effective prescription treatment that has no known published data documenting human mite resistance to the active compound, spinosad.³

Visit natroba.com

References: 1. IQVIA May 28, 2021. 2. Data on file, ParaPRO, LLC. 3. Natroba Prescribing Information. 4. JAAD clinical article (full citation TBD). 5. Khalil S et al. *PLoS Negl Trop Dis*. 2017;11(11): e0005920.

IMPORTANT SAFETY INFORMATION

INDICATION

Natroba™ Topical Suspension is a scabicide indicated for the topical treatment of scabies infestations in adult and pediatric patients four (4) years of age and older.

ADJUNCTIVE MEASURES

Natroba™ Topical Suspension should be used in the context of an overall lice management program:

- Wash in hot water or dry-clean any bedding, clothing and towels used by anyone having scabies.
- If any member of a household presents with scabies, all household members should be treated with Natroba™ Topical Suspension.



IMPORTANT SAFETY INFORMATION

Natroba™ Topical Suspension contains benzyl alcohol and is not recommended for use in pediatric patients below the age of 4 years. The safety and effectiveness of Natroba™ Topical Suspension have not been established in pediatric patients less than 4 years of age with scabies infestation. Most common adverse events were: application site irritation (3%)* and dry skin (2%).

*Application site irritation also includes application site pain and burning sensation.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

See pocket for Full Prescribing and Patient Information.